

CALLIOPSIS

Coreopsis tinctoria Nutt.

plant symbol = COTI3

Contributed by: USDA NRCS Plant Materials Program



© Jim Stasz
MD National Capital Park &
Planning Commission
@USDA NRCS PLANTS

Alternate Names

Plains coreopsis, golden tickseed

Uses

This plant is used mainly for landscape beautification. It has potential for use in cultivated, garden situations, in naturalized prairie or meadow plantings, and along roadsides.

Status

Please consult the PLANTS Web site and your State Department of Natural Resources for this plant's current status (e.g. threatened or endangered species, state noxious status, and wetland indicator values).

Description

Calliopsis is an annual forb which usually germinates in late summer or fall and overwinters as a rosette (round, low growing group of leaves). The stems begin growing upward and branching in spring, reaching a height of up to 4 feet. The opposite leaves are deeply divided, with the upper leaf segments being very narrow. The flowering heads are numerous, 1-2 inches in diameter, and are yellow with a red-brown center.

Adaptation

Calliopsis is adapted to many soil types. It grows best on a well-drained soil, but will not tolerate a very dry site. Natural stands are generally found on bottomland areas with ample moisture. It grows best in full sun, but will tolerate light shade.

Calliopsis is distributed throughout the majority of the United States. For a current distribution map, please consult the Plant Profile page for this species on the PLANTS Website.

Establishment

A firm seedbed is required. Calliopsis germinates best on a clean tilled site that has been firmed with a roller or finishing harrow before planting. Seed can also be planted into a closely mowed, chemically-killed, or burned sod area with a light disking or harrowing that scratches the soil surface. A layer of plant residue on the soil surface can interfere with seed germination. Broadcast or shallowly drill 1-2 grams per 100 square feet (1-2 lb/acre). Seed should be placed close to the soil surface. Cultipacking after planting will ensure good seed to soil contact. Seed will germinate soon after planting and remain as a rosette throughout the winter.

Management

Plant growth and seed production are greatly improved by fertilization. Apply fertilizer according to soil test recommendations. If not available, a rate of 3.5-5.5 oz per 100 square feet (100-150 lb/acre) of 13-13-13 should be applied in the spring before flowering. Stands will reseed prolifically for several years, but will gradually decline without soil disturbance. Every two to three years, the site should be disked to control perennial weeds and promote calliopsis germination. If necessary, plants can be mowed in the spring before stem elongation begins. Stands that are not disked should be mowed in late summer, and a late fall mowing is also recommended.

Prepared By & Species Coordinator:

USDA, Natural Resources Conservation Service
Jamie L. Whitten Plant Materials Center
Route 3, Box 215-A
Coffeeville, MS 38922-9263

01Feb2002 JLK

For more information about this and other plants, please contact your local NRCS field office or Conservation District, and visit the PLANTS <<http://plants.usda.gov>> and Plant Materials Program Web sites <<http://Plant-Materials.nrcs.usda.gov>>.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.