

MARYLAND MEADOW BEAUTY

Rhexia mariana L.

plant symbol = RHMA

Contributed by: USDA NRCS Plant Materials Program



Robert H. Mohlenbrock
USDA NRCS 1989
Midwestern Wetland Flora
@ USDA NRCS PLANTS

Uses

This plant is used mainly for landscape beautification. It has potential for use in cultivated, garden situations, in naturalized prairie or meadow plantings, and along roadsides.

Status

Please consult the PLANTS Web site and your State Department of Natural Resources for this plant's current status (e.g. threatened or endangered species, state noxious status, and wetland indicator values).

Description

An upright (up to 2.5 feet), hairy perennial that frequently spreads to form large colonies. The opposite leaves are elongated, three-nerved, with toothed margins. The pink to lavender or occasionally white flowers are produced from May to October in small clusters near the top of the plant. Each flower is 1-2 inches in diameter and contains 4 petals; the yellow to orange anthers in the center are elongated and curved downwards.

The tiny tan to brownish seeds are held in a reddish urn-shaped capsule.

Adaptation and Distribution

Meadow beauty grows best in a sunny exposure. It is adapted to heavy textured and to highly organic soils. Natural stands are usually found on moist to wet sites such as in ditches, marshes, and wet meadows.

Maryland meadow beauty is distributed primarily throughout the East and lower Midwest. For a current distribution map, please consult the Plant Profile page for this species on the PLANTS Website.

Establishment

A clean, firm seedbed is essential. The site should be treated with a herbicide to control vegetation, tilled, cultipacked once or twice, and allowed to settle thoroughly before sowing.

Broadcast 0.5-1 gram seed per 100 square feet (0.5-1 lb/acre). The seed can be mixed with sand or rice hulls to increase volume so that it will be easier to spread uniformly over the planting site. The seed must remain on the soil surface because it is easily smothered when buried in the soil. Heavy rains after planting may cause the seed to be buried or washed from the site.

Management

Apply fertilizer according to soil test recommendations. If not available, a rate of 3.5-5.5 oz per 100 square feet (100-150 lb/acre) of 13-13-13 should be applied after the seedlings are established and annually thereafter. If necessary, stands can be mowed in the spring before stem elongation begins. Later mowings should be delayed until the plants have set seed.

Prepared By & Species Coordinator:

USDA NRCS Plant Materials Program

05Feb2002.JLK

For more information about this and other plants, please contact your local NRCS field office or Conservation District, and visit the PLANTS <<http://plants.usda.gov>> and Plant Materials Program Web sites <<http://Plant-Materials.nrcs.usda.gov>>.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.