

SOUTHERN ARROWWOOD

Viburnum dentatum L.
 plant symbol = VIDE

Contributed By: USDA, NRCS, National Plant Data Center & the Biota of North America Program



© Hugh Wilson
 Vascular Plant Image Gallery
 Texas A&M University, Bioinformatics Working Group

Alternate Names

Arrow-wood viburnum, southern arrow-wood, roughish arrow-wood. (Arrowwood is frequently hyphenated: arrow-wood.) The common name refers to the Native American use of the straight young stems as arrow shafts.

Uses

Landscape: *V. dentatum* is an adaptable native, multi-stemmed shrub. Creamy white flowers, dark blue berries and colorful fall foliage make southern arrowwood an attractive landscape plant. It suckers freely from the base. It can be used for borders or screens or as mass plantings and groupings to attract birds, which eat the fruit.

Medicinal: *Viburnum* species have been used for numerous medicinal purposes (Alternative Medicine Foundation: HerbMed, 2000--for notes and internet links on medicinal use and other health related topics for *Viburnum* species).

Status

Please consult the PLANTS Web site and your State Department of Natural Resources for this plant's current status (e.g. threatened or endangered species, state noxious status, and wetland indicator values).

Description

Southern arrowwood is a native shrub growing 3-9 feet tall and spreading sometimes up to 8 feet. The plant's arching branches form an overall rounded crown; twigs are slender, ridged and angled. Its leaves are deciduous, opposite, simple, oval to oblong, obovate (inversely egg-shaped), or elliptic in shape and 1½ - 4 inches in length, with coarsely, but regularly, toothed margins. Leaves are shiny dark green above and paler beneath, sparsely stellate (with hairs in small star-like tufts) on undersides and petioles. Foliage turns yellow to red or reddish-purple in late fall. Small white flowers are borne in 2 to 4-inch flat-topped clusters in May to early June. The ¼ inch berry-like drupes are bluish-black and attractive to wildlife. Fruiting occurs from August - November.

Localized variations of southern arrowwood occur over the geographic range of the species. Most common differences between the variants are in the shape and size of leaves, the type and placement of pubescence (hairs) on the leaf underside and petioles, and the region of occurrence. Some variations of southern arrowwood include *V. dentatum* var. *dentatum*; *V. dentatum* var. *scabrellum* Torr. & Gray (= *V. scabrellum* (Torr. & Gray) Chapm.) and *V. dentatum* var. *venosum* (Britt.) Gleason.

Adaptation and Distribution

Southern arrowwood is found natively in open woods and margins, and along streambanks. It prefers loamy, neutral to acid soil with ample moisture, but is adaptable to a range of conditions from dry to fairly wet soil. Plants are salt-tolerant in New England coastal areas. They can grow in generally drier conditions than *V. acerifolium*. They most commonly occur in partial shade but can be grown in full sun.

Southern arrowwood is distributed throughout the East. For a current distribution map, please consult the Plant Profile page for this species on the PLANTS Website.

Establishment

Southern arrowwood plants are propagated by seed or vegetatively. Seed bearing age begins at approximately 3-4 years. Depending on region of collection, the seeds of *V. dentatum* may have a cold requirement for breaking embryo dormancy. Vegetative reproduction is through short rhizomes and sprouts from the root crown or softwood cuttings. Southern arrowwood has a fibrous root system and is easily transplanted.

Management

V. dentatum is free from serious problems of disease or most insect pests (see, however, discussion on viburnum leaf beetle below). Occasional pruning is helpful to rejuvenate and shape the plant. Prune off basal suckers to restrict spreading if necessary.

Pests and Potential Problems

The viburnum leaf beetle (*Pyrrhalta viburni*) was introduced from Europe and Asia to North America around 1947. It became a problem in Canada in 1978 and has now moved to the northeastern United States where it is a concern in urban landscapes and nurseries. *Viburnum dentatum* leaves can be damaged or skeletonized by the beetle adults and larvae, though *Viburnum opulus* (European cranberrybush viburnum) appears to be the insect's preferred host and is most seriously affected. The beetle larvae hatch in early May, feed for about 4-5 weeks then pupate in the soil. Adults emerge by mid-July, feed, mate, and females lay over-wintering eggs in a straight line on viburnum twigs. If found, the eggs should be pruned out and destroyed before hatching. Chemical control is best applied to young larvae, which feed on both upper and lower leaf surfaces.

Cultivars, Improved, and Selected Materials (and area of origin)

Several cultivars of *V. dentatum* are available.

References

See the plant guide for this species to obtain a complete list of text references.

Prepared By:

Guy Nesom

Bonap, North Carolina Botanical Garden,
University of North Carolina, Chapel Hill, North
Carolina

Kathy Davis

USDA-NRCS National Plant Materials Center,
Beltsville, Maryland

Species Coordinator:

Lincoln Moore

USDA, NRCS, National Plant Data Center, Baton
Rouge, Louisiana

For more information about this and other plants, please contact your local NRCS field office or Conservation District, and visit the PLANTS <http://plants.usda.gov> and Plant Materials Program Web sites <http://Plant-Materials.nrcs.usda.gov>.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.