

## AMUR MAPLE

*Acer ginnala* Maxim.  
plant symbol = ACGI

Contributed By: USDA, NRCS, National Plant Data Center



from Conservation Trees and Shrubs for Montana  
USDA NRCS Montana State Office

### Alternative Name

Siberian maple

### Uses

*Ethnobotanic:* The young leaves were used as a tea substitute (Kunkel 1984). Black, blue, and brown dyes were obtained and dried from the leaves.

*Landscaping & Wildlife:* The main ornamental value of Amur maple is its red fall color and fruit. This is an excellent, low growing tree for small yards. It is sometimes used for hedges or screens. It has a fair rating for wildlife.

*Agroforestry:* *Acer ginnala* is used in tree strips for windbreaks. They are planted and managed to protect livestock, enhance production, and control soil erosion. Windbreaks can help communities with harsh winter conditions better handle the impact of winter storms and reduce home heating costs during the winter months.

### Status

Introduced into the U.S. Please consult the PLANTS Web site and your State Department of Natural Resources for this plant's current status, such as, state noxious status and wetland indicator values.

### Description

*General:* Maple family (Aceraceae). Amur maple is an introduced, deciduous large shrub or small tree. It can be grown as a multistemmed clump or trained into a small tree with a single trunk. It can also be sheared into a hedge. The leaves are simple, opposite; eight to ten centimeters long, and coarsely toothed. The fragrant, creamy white flowers appear with the new foliage in April and May (Dirr 1997). The fruit samaras are 0.75 to 1 inch long, held in small panicles and are red to brown in color. The bark is smooth and gray on young branches and grayish brown on older branches.

*Distribution:* Amur maple is native to Manchuria, north China and Japan and is cultivated in gardens and parks (McMinn & Maino 1951). For current distribution, please consult the Plant profile page for this species on the PLANTS Web site.

### Adaptation

*Acer ginnala* grows best in moist well-drained soil, but can tolerate a wide variety of soils, poor soil fertility, and are pH adaptable. This species displays excellent tolerance to dry and alkaline soils (Dirr 1997). It is reasonably drought tolerant. It will tolerate shade, but develops a better fall color if grown in full sun. *Acer ginnala* grows best in colder climates with cool summers. It can be grown in hotter areas if care is taken to prevent dehydration.

### Establishment

*Propagation from Seed:* Pre-soak the stored seed for twenty-four hours and then stratify for one to four months at 1-8°C. Seeds can be harvested when they are fully developed but before they have dried and produced any germination inhibitors. Sow immediately in a seedbed or open frame. Spring sown seeds may not germinate for another year. Transfer to a nursery bed in the first spring (Heuser 1997). If the seeds are harvested too soon they will produce very weak plants or no plants at all (McMillan 1985).

*Propagation from Cuttings:* Cuttings of young shoots should be done in June or July. The cuttings should consist of two to three pairs of leaves and one pair of buds on the base. Place cuttings in plastic, bag and seal to prevent moisture loss. They must not be allowed to wilt. Trim the cuttings below the lowest

node to remove the lower leaves leaving three or four at the tip. A rooting hormone may be applied to improve rooting before planting. Insert the cuttings in the rooting medium up to half their length so the leaves don't touch each other. The cuttings should root in two to three weeks, after which they can be potted (Heuser1997).

### Management

Containerized trees should not be planted in their permanent position until they are twenty centimeters or taller. Amur maple should be pruned in the winter or early spring to help develop and maintain a good single trunk tree form. Amur maple is usually pest free; however, sometimes, spraying is necessary to control aphids.

### Cultivars, Improved and Selected Materials (and area of origin)

'Red Fruit', 'Bailey Compact', 'Flame', 'Compactum' and 'Durand Dwarf' are cultivars of *Acer ginnala*. 'Red Fruit' is a collective term for types whose fruit color is brilliant red (Dirr 1990). 'Bailey Compact' has a compact shrubby form, growing between eight to twelve inches high. 'Flame' is a dense shrub or small tree with red fruits and fiery red fall color (Dirr 1990). 'Compactum' is dense and compact and shows vigorous growth reaching between five to six inches. 'Durand Dwarf' is a shrubby type, with branches more dense than 'Compactum', will grow three to five inches high.

Consult your local nurseries to choose the right cultivar for your specific landscape.

### References

Barnes, B.V. & W.H. Wagner, Jr. 1981. *Michigan trees*. The University of Michigan Press, Ann Arbor, Michigan.

Dirr, M.A. 1997 *Dirr's hardy trees and shrubs: an illustrated encyclopedia*. Timber Press, Portland, Oregon.

Dirr, M.A. 1990. *Manual of woody landscape plants: their identification, ornamental characteristics, culture, propagation, and uses*. 4<sup>th</sup> ed. Stipes Publishing Co., Champaign, Illinois.

Dirr, M.A. & C.W. Heuser, Jr. 1987. *The reference manual of woody plant propagation: from seed to tissue culture*. Varsity Press, Athens, Georgia.

Farrar, J.L. 1995. *Trees of the Northern United States and Canada*. Iowa State University Press, Ames, Iowa.

Heuser, C.W. 1997. *The complete book of plant propagation*. The Taunton Press, Newtown, Connecticut.

McMillan, B.P. 1985. *Hardy woody plants of North America*. Grower Books.

McMinn, H.E. 1951. *An illustrated manual of California shrubs*. University of California Press, Berkeley & Los Angeles, California.

Kunkel, G. 1984. *Plants for human consumption*. Koeltz Scientific Books.

Rosendahl, C.O. 1955. *Trees & shrubs of the upper Midwest*. University of Minnesota Press, Minneapolis, Minnesota.

Taylor, N. 1965. *The guide to garden shrubs and trees*. Houghton Mifflin Company, Boston, Massachusetts.

USDA, NRCS 2000. *Conservation trees and shrubs for Montana*. Custer County Soil Conservation District. Accessed: 10jan02.  
<<http://www.mt.nrcs.usda.gov/pas/forestry/maple.html>>

Wyman, D. 1965. *Trees for American gardens*. The MacMillan Company, New York, New York .

### Prepared By

*Jammie Favorite*

formerly USDA, NRCS, National Plant Data Center  
Baton Rouge, Louisiana

### Species Coordinator

*Lincoln M. Moore*

USDA, NRCS, National Plant Data Center  
Baton Rouge, Louisiana

Edited: 10jan02 jsp

For more information about this and other plants, please contact your local NRCS field office or Conservation District, and visit the PLANTS Web site <<http://plants.usda.gov>>.

*The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).*

*To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.*