

DOWNY SERVICEBERRY

Amelanchier arborea (Michx.
f.) Fern.
plant symbol = AMAR3

Contributed By: USDA, NRCS, National Plant Data Center & Biota of North America Program



© William S. Justice
Botany Dept., NMNH, Smithsonian Institution
@ PLANTS

Alternate common names

Sarvis-berry, shadblow, shadbush, juneberry, sugarplum, Indian cherry

Uses

Trees of downy serviceberry are generally not large enough for sawtimber but they have been used for pulpwood. The wood is extremely heavy and hard and is occasionally made into tool handles. It was prized by Cree Indians for making arrows.

At least 40 bird species (for example, mockingbirds, cardinals, cedar waxwings, towhees, Baltimore orioles) eat the fruit of *Amelanchier* species.

Mammals that either eat the fruit or browse the twigs and leaves of downy serviceberry include squirrels, rabbits, chipmunks, mice, voles, foxes, black bears, deer, and elk. The fruits taste similar to blueberry – they are eaten fresh or cooked in pastries or puddings.

The trees are used as ornamentals and many cultivars have been selected for variation in growth habit, flower size and color, and leaf color. The fall foliage blends orange and gold with red and green. It grows

in partial shade to full sun, preferring moist but well-drained soil but will also grow in dry sites.

Status

Please consult the PLANTS Web site and your State Department of Natural Resources for this plant's current status, such as, state status and wetland indicator values.

Description

Rose Family (Rosaceae). Native shrubs or small trees to 10 meters tall, with a narrow, rounded crown, the twigs often red-brown to purplish, becoming gray; bark smooth, grayish, "striped" with vertical fissures and very ornamental. Leaves: deciduous, alternate, simple, oval to oblong, 5-13 cm long, glabrous above, pubescent and paler beneath, the base rounded or heart-shaped, acute or acuminate at the tip, with finely toothed margins. Flowers: 3-15 in elongate clusters at the branch tips, before the leaves appear; petals 5, white, 10-14 mm long and strap-like. Fruits 6-12 mm wide, on long stalks, red-purple at maturity; seed 5-10 per fruit. The common name: in some regions, the flowers are gathered for church services, hence serviceberry or sarvis-berry; or "service" from "sarvis," in turn a modification of the older name "*Sorbus*," a closely related genus.

Variation within the species: Three varieties have been recognized: var. *alabamensis* (Britt.) G.N. Jones; var. *arborea*; and var. *austromontana* (Ashe) Ahles.

Distribution

Downy serviceberry is widespread in the eastern US and southeastern Canada (New Brunswick and southern Newfoundland to Quebec and Ontario); south to the northern tip of the Florida Panhandle and west to Alabama, southern Mississippi, Louisiana and Texas (rare), Oklahoma, Kansas, Nebraska, and Minnesota.

Adaptation

Downy serviceberry grows in a variety of habitats – swampy lowlands, dry woods, sandy bluffs, rocky ridges, forest edges, and open woodlands and fields. It is a late successional to climax species in mixed-hardwood forests of the central U.S., commonly as an understory species. In the southern Appalachians, downy serviceberry grows in red spruce-Fraser fir forests at elevations of 1500-2000 meters with yellow birch, mountain ash, elderberry, and hobblebush. Flowering (March-)April-May, among the first of the

early spring trees and shrubs to bloom; fruiting June-August.

Establishment

Downy serviceberry regenerates mainly by seed, but it also sprouts from the roots. Seeds are dispersed by birds and mammals; scarification of the seeds after ingestion by birds is important for germination. Seeds can be sown after 2-6 months of cold stratification, but they will not usually germinate until after the second spring.

Management

Fire top-kills downy serviceberry, but it can sprout from root crowns and stumps following fire. A significant portion of the post-fire reestablishment is from seed dispersed from off-site by birds and mammals. Following wildfire in a spruce-fir forest of Appalachia, downy serviceberry was present in stands after 30 years but was less than 1% of the total basal area. Gypsy moth larvae (*Lymantria dispar*) feed selectively on downy serviceberry.

Cultivars, Improved and Selected Materials (and area of origin)

Please check the Vendor Database, expected to be on-line through the PLANTS Web site in 2001 by clicking on Plant Materials. Natural, fertile hybrids occur between downy serviceberry and *A. bartramiana*, *A. canadensis*, *A. humilis*, and *A. laevis*. Some cultivars are selections from *A. X grandiflora*, the hybrid of *A. arborea* and *A. laevis*. Many individuals within *Amelanchier* arise through hybridization and species boundaries are often not clear.

References

Campbell, C.S., M.F. Wojciechowski, B.G. Baldwin, L.A. Alice, & M.J. Donoghue 1997. *Persistent nuclear ribosomal DNA sequence polymorphism in the Amelanchier agamic complex (Rosaceae)*. Molecular Biol. Evol. 14:81-90.

Campbell C.S., W.A. Wright, T.F. Vining, & W.A. Halteman 1997. *Morphological variation in sexual and agamosperous Amelanchier (Rosaceae)*. Canad. J. Bot. 75:1166-1173.

Campbell, C.S., C.W. Greene, & S.E. Bergquist 1987. *Apomixis and sexuality in three species of Amelanchier, shadbush (Rosaceae, Maloideae)*. Amer. J. Bot. 74:321-328.

Campbell, C.S. & W.A. Wright 1996. *Apomixis, hybridization, and taxonomic complexity in eastern*

North American Amelanchier (Rosaceae). Folia Geobot. Phytotax. 31:345-354.

Dibble, A.C. 1995. *Conservation biology of shadbush, Amelanchier (Rosaceae): Evidence from systematics, population structure and reproductive ecology (Amelanchier nantucketensis, Amelanchier stolonifera, Amelanchier obovalis, Amelanchier lucida)*. Ph.D. diss., Univ. of Maine, Orono, Maine.

Harris, R.E. 1970. *The genus Amelanchier*. J. Roy. Hort. Soc. 95:116-118.

Iverson, R. 1981. *Amelanchiers*. Amer. Horticult. 60(10):26-27, 36-37.

Landry, P. 1976. *Taxonomy and distribution of Amelanchier arborea (Michx. f.) Fern. in Quebec and through the Maritimes*. Naturaliste Canad. 103:377-385.

McDaniel, J.C. 1974. *Nomenclature: perennial problem in Amelanchier arborea*. Amer. Nurseryman 139(7):66-68.

Overath, R.D. & J.L. Hamrick 1998. *Allozyme diversity in Amelanchier arborea and A. laevis (Rosaceae)*. Rhodora 100:276-292.

Snyder, S.A. 1992. *Amelanchier arborea*. IN: W.C. Fischer (compiler). *The fire effects information system* [Database]. USDA, Forest Service, Intermountain Research Station, Intermountain Fire Sciences Laboratory, Missoula, Montana. <<http://www.fs.fed.us/database/feis/>>

Robinson, W.A. 1986. *Effect of fruit ingestion on Amelanchier seed germination*. Bull. Torrey Bot. Club 113:131-134.

Weaver, R.E. 1974. *The shadbushes*. Arnoldia (Jamaica Plain) 34:22-31.

Prepared By

Guy Nesom
BONAP, North Carolina Botanical Garden,
University of North Carolina, Chapel Hill, North Carolina

Species Coordinator

Lincoln Moore
USDA, NRCS, National Plant Data Center, Baton Rouge, Louisiana

Edited: 17jan01 jsp

For more information about this and other plants, please contact your local NRCS field office or Conservation District, and visit the PLANTS <<http://plants.usda.gov>> and Plant Materials Program Web sites <<http://Plant-Materials.nrcs.usda.gov>>.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.