

CROSSVINE

Bignonia capreolata L.

Plant symbol = BICA

Contributed by: USDA NRCS National Plant Data Center



J.R. Manhart. 2004.
Texas A&M University Herbarium.

Alternate Names

Anisostichus capreolata, *Anisostichus crucigera*, *Bignonia capreolata* forma *lutea*, *Bignonia capreolata* var. *atrosanguinea*, *Doxantha capreolata*, quartervine, trumpet flower.

Uses

Ethnobotanic: Native Americans used crossvine as a remedy for numerous health ailments. An infusion of leaves was used to purify blood. Decoctions of leaves were used for rheumatism. Decoctions of mashed bark were used to alleviate edema and headaches. Individuals with diphtheria gargled a mashed root infusion.

Ornamental: The showy, fragrant flowers and unique leaves of crossvine make this plant appropriate for some gardening and landscaping needs. It is often used as a cover for fences, arbors, walls, pillars or large trellises and as a groundcover. It is often recommended as a native ornamental alternative to non-native climbing vines.

Wildlife: The tubular flowers and large quantities of nectar produced by crossvine are attractants for hummingbirds and butterflies.

Legal Status

Please consult the PLANTS Web site (<http://plants.usda.gov>) and your State Department of Natural Resources for this plant's current status (e.g. threatened or endangered species, state noxious status, and wetland indicator values).

Weediness

Crossvine can spread aggressively. This plant may become weedy or invasive in some regions or habitats and may displace desirable vegetation if not properly managed. Please consult with your local NRCS Field Office, Cooperative Extension Service office, or state natural resource or agriculture department regarding its status and use. Weed information is also available from the PLANTS Web site at plants.usda.gov.

Description

General: Bignonia family (Bignoniaceae). Crossvine is a perennial, woody, semi-evergreen vine that can grow to be 17 m long. Tubers can grow to 10 cm in diameter. The stem is squarish and reddish-purple. Leaves are opposite, pinnately compound, and reduced to two basal leaflets and a branched tendril between them. Leaflets are 6-15 cm long, 2-7 cm wide and have smooth edges (entire), narrowly tapered tips (acuminate), and a notch at the base that makes them heart-shaped (cordate). The foliage turns from a lustrous green in the growing season to a reddish purple in the winter. Buds are reddish-purple and less than 1 cm long. Flowers are bell-shaped with orange on the outside and yellow inside, have 5 irregular lobes, and are 4-5 cm long. The fruit is a multi-seeded, flat capsule that is up to 15 cm long. Crossvine blooms from mid-March to mid-June and sets fruit in late summer.

Crossvine gets its name from the cross-shaped pith, as seen from a transverse section of the stem. Other key characteristics of crossvine include its opposite compound leaves with two leaflets and branched tendrils.

Distribution: Crossvine is native to the United States in the northeastern, north central, south central, and southeastern regions. It is found from southern Ontario, south to Florida, west to Texas and Illinois. For current distribution, please consult the Plant Profile page for this species on the PLANTS Web site (<http://plants.usda.gov>).

Habitat: Crossvine is found in rich forests and swamps and along roadsides and fencerows.

Adaptation

The USDA hardiness zones for crossvine are 5-9. Although it is semi-evergreen, it may lose its leaves during especially harsh winters in zones 5 and 6.

Crossvine prefers full sun and moist, acidic, well-drained soils for best flowering. It is adaptable to other soil conditions, including poorly drained soils, once established and it can do well in low light levels. It is drought tolerant.

Management

If not properly managed, crossvine can spread quickly by root suckers and become problematic. Pruning will reduce aggressive growth if spread is not desired.

Seeds and Plant Production

Crossvine can be propagated by seeds or cuttings. Pretreatment of seeds is not necessary. Although a 90% germination rate can be expected, seedling survival and growth may be low. Seedlings may need to harden in winter temperatures before outplanting.

The species can be rooted at any time of the year but June or July cuttings are generally recommended. Crossvine cuttings root easily, but may take six to eight weeks.

Cultivars, Improved, and Selected Materials (and area of origin)

The NRCS Plant Materials Program has not released any cultivars of crossvine for conservation use.

Ornamental cultivars of *Bignonia capreolata* include 'Atrosanguinea,' 'Dragon Lady,' 'Helen Fredel,' 'Jekyll,' 'Shalimar Red,' and 'Tangerine Beauty.' These cultivars have been bred for flower color, flower and leaf size, and winter hardiness.

Control

Please contact your local agricultural extension specialist or county weed specialist to learn what works best in your area and how to use it safely. Always read label and safety instructions for each control method. Trade names and control measures appear in this document only to provide specific information. The USDA NRCS does not guarantee or warranty the products and control methods named, and other products may be equally effective.

References

Agricultural Research Center. 2004. *GRIN taxonomy* (<http://www.ars-grin.gov/cgi-bin/npgs/html/index>, 22 April 2004). USDA, Beltsville.

Creech, D. 2004. *Plants page, SFA Mast Arboretum* (<http://arboretum.sfasu.edu/plants/index.htm>, 22 April 2004). Stephen F. Austin State University, Nacogdoches.

Dirr, M.A. 2004. *Horticulture 3710 plant list: vines* (<http://www.nobleplants.com/classnotes/spring/springprofiles/vines.htm>, 22 April 2004). Noble Plants, Athens.

Floridata. 2004. *Plant encyclopedia* (<http://floridata.com>, 22 April 2004). Floridata Marketplace, Tallahassee.

Johnson, F.L. and B.W. Hoagland. *Catalog of the woody plants of Oklahoma* (<http://www.biosurvey.ou.edu/shrub/cover.htm>, 22 April 2004). Oklahoma University, Norman.

Kemper Center for Home Gardening. 2004. *Kemper web, PlantFinder* (<http://ridgwaydb.mobot.org/kemperweb/plantfinder/Alpha.asp>, 22 April 2004). Missouri Botanical Garden, St. Louis.

Lindstrom, J.T. 2002. *Bignonia capreolata* (http://www.uark.edu/campus-resources/cotinus/plants3_html/bigncapr.html, 22 April 2004). University of Arkansas, Fayetteville.

Manhart, J.R. 2004. *Texas A & M University Herbarium digital flora of Texas* (<http://www.csdl.tamu.edu/FLORA/gallery1.htm>, 22 April 2004). Texas A & M University, College Station.

Reed, D. 2002. *Wildflowers of the Southeastern United States* (<http://2bnthewild.com/>).

Stine, M. 2003. *Louisiana ecosystems and plant identification*, (<http://mrstreamer.lsu.edu/ecosystems/webtour/>, 22 April 2004). LSU AgCenter, Baton Rouge.

Prepared By:

Sarah Wennerberg
USDA NRCS National Plant Data Center
Baton Rouge, Louisiana

Species Coordinator:

Mark Skinner

USDA NRCS National Plant Data Center

Baton Rouge, Louisiana

Edited: 6May2004 sbw; 21Oct2004 rln

For more information about this and other plants, please contact your local NRCS field office or Conservation District, and visit the PLANTS Web site <<http://plants.usda.gov>> or the Plant Materials Program Web site <<http://Plant-Materials.nrcs.usda.gov>>

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.