

## MAPLE-LEAF VIBURNUM

*Viburnum acerifolium* L.

plant symbol = VIAC

Contributed By: USDA, NRCS, National Plant Data Center & the Biota of North America Program

### Alternate common names



Botany Dept., NMNH, Smithsonian Institution  
@ PLANTS

Dockmackie, guelder-rose, maple-leaved arrow-wood, possum-haw, squash-berry

### Uses

The fruits of maple-leaf viburnum are eaten by deer, rabbits, mice, skunks, ruffed grouse, ring-necked pheasants, wild turkeys, and many species of songbirds. The twigs, bark, and leaves are eaten by deer, moose, rabbits, and beavers. The relatively low-growing plants provide good nesting and escape cover for birds and small mammals.

Maple-leaf viburnum has long been cultivated for its attractive summer flowers and foliage; then the autumn leaves turn rose-purple and contrast with the mature dark fruits. The plants will thrive in moist soils and a range of light conditions but they are a good choice for dry soils in deep shade. They can be used along forest edges, streamsides, and lakeshores.

### Status

Please consult the PLANTS Web site and your State Department of Natural Resources for this plant's current status, such as, state noxious status and wetland indicator values.

### Description

*General:* Honeysuckle family (Caprifoliaceae). Native shrubs 0.5-2 m high, rhizomatous and forming

thickets; twigs pubescent. Leaves deciduous, ovate or orbicular, mostly deeply 3-lobed, 5-12 cm broad, coarsely toothed, lower surface minutely black-dotted, nearly glabrous to thinly pubescent with stellate hairs. Flowers white, bisexual, 4-6 mm wide, in upright, flat-topped clusters 4-7 cm wide. Fruit berry-like (drupes), 6-8 mm long, nearly black, with a single stone.

### Variation within the species:

No varieties are currently formally recognized within *V. acerifolium* – previous named varieties within the species have described vaguely discernible and widely overlapping geographic trends of morphological variation.

*V. acerifolium* var. *acerifolium*

*V. acerifolium* var. *densiflorum* (Chapm.) McAtee

*V. acerifolium* var. *glabrescens* Rehd.

*V. acerifolium* var. *ovatum* (Rehd.) McAtee

*Distribution:* Widely distributed in eastern North America, New Brunswick (rare), Quebec, and Ontario south through Wisconsin, Illinois, and Arkansas to Florida and eastward into east Texas. For current distribution, please consult the Plant Profile page for this species on the PLANTS Web site.

### Adaptation

Maple-leaf viburnum occurs in upland forests, hillsides, and ravine slopes. It grows best in well-drained, moist soils and is tolerant of acid soils. It requires partial shading for optimum growth and development and occurs primarily in mid- to late-seral communities. It is a common understory species in beech-maple forests in the northeastern and midwestern United States; along the Gulf coastal plain, it is found in rich deciduous woods, often with white oak. Flowering May-August; fruiting July-October.

### Establishment

Maple-leaf viburnum begins seed production at about 2 years of age and produces abundant fruit every year. Most seeds have an impermeable seedcoat and embryo dormancy that requires a warm-cold stratification sequence to be broken. Vegetative reproduction through rhizomes is extensive.

### Management

Low- to moderate-severity fires top-kill maple-leaf viburnum. It apparently survives fire by sprouting from underground rhizomes, but these are shallow

and easily damaged and the species decreases with exposure to repeated fires.

*Viburnum leaf beetle*. The viburnum leaf beetle (*Pyrrhalta viburni*), native to Europe and Asia, was first encountered in North America in 1947, perhaps arriving earlier from Europe on nursery plants. It received little notice until 1978, when it caused severe defoliation of ornamental viburnums in Ontario and Quebec. It has now reached western New York and Maine and become a concern in urban landscapes and nurseries.

The adult and the larva “skeletonize” leaves by feeding on the leaves between the midrib and larger veins. Plants which have been defoliated for 2-3 consecutive years may be killed. The preferred host is *Viburnum opulus* and its selections; lesser damage is caused to *V. lantana* and *V. rafinesquianum*, *V. dentatum*, *V. acerifolium*, and *V. lentago*. Other species, particularly *V. rhytidophyllum* and *V. carlesii*, are relatively unaffected.

The entire life cycle of the viburnum leaf beetle takes about 8-10 weeks. Larvae hatch in early May and feed on the viburnum leaves throughout the larval period, which lasts 4-5 weeks. The larvae pupate in the soil. The adults (4.5-6.5 mm long, brown) appear by mid-July and continue eating the leaves, then mate and lay overwintering eggs on the twigs. Egg-laying holes are in a straight line on the underside of the current season's growth.

Chemical control of the viburnum leaf beetle is best applied to young larvae, because adults will fly away or drop to the ground if disturbed. If over-wintering egg sites are found, affected wood should be pruned and destroyed before the eggs hatch. Examine upper and lower leaf surfaces for feeding larvae. Potential biological control mechanisms are being studied.

#### **Cultivars, Improved and Selected Materials (and area of origin)**

Please check the Vendor Database, expected to be on-line through the PLANTS Web site in 2001 by clicking on Plant Materials. These plant materials are readily available from commercial sources.

#### **References**

Coladonato, M. 1993. *Viburnum acerifolium*. IN: W.C. Fischer (compiler). *The fire effects information system* [Data base]. U.S.D.A, Forest Service, Intermountain Research Station, Intermountain Fire Sciences Laboratory. Missoula, Montana.  
<<http://www.fs.fed.us/database/feis/>>

Donoghue, M. 1980. *Flowering times in Viburnum*. *Arnoldia* 40:2-22;

Donoghue, M. 1983. *A preliminary analysis of phylogenetic relationships in Viburnum (Caprifoliaceae s.l.)*. *Syst. Bot.* 8:45-58.

Egolf, D.R. 1962. *A cytological study of the genus Viburnum*. *J. Arnold Arb.* 43:132-172.

Friedlander, B.P., Jr. 1999. *Voracious viburnum leaf beetle munches into Ithaca area*. *Cornell Chronicle*. <[http://www.news.cornell.edu/Chronicles/6.17.99/leaf\\_beetle.html](http://www.news.cornell.edu/Chronicles/6.17.99/leaf_beetle.html)>

Hauser, E.J.P. 1965. *Characteristics and distribution of Viburnum (Caprifoliaceae) in Georgia*. *Bull. Ga. Acad. Sci.* 23:(11 pages).

Kessel, C. 2000. *Viburnum leaf beetle, Pyrrhalta viburni (Paykull), in the nursery and landscape*. Ontario Ministry of Agriculture, Food and Rural Affairs, Ontario, Canada. Web site. <<http://www.gov.on.ca/OMAFRA/english/crops/facts/vlb.htm>>

McAtee, W.L. 1956. *A review of the Nearctic Viburnum*. Published by the author, Chapel Hill, North Carolina.

Thomson J. & O. Thomson 1981. *Leaf shapes of Viburnum acerifolium*. *Bull. Bot. Club Wisconsin* 12:4-5.

#### **Prepared By**

*Guy Nesom*  
BONAP, North Carolina Botanical Garden,  
University of North Carolina, Chapel Hill, North Carolina

#### **Species Coordinator**

*Lincoln Moore*  
USDA, NRCS, National Plant Data Center, Baton Rouge, Louisiana

Edited 17jan01 jsp

For more information about this and other plants, please contact your local NRCS field office or Conservation District, and visit the PLANTS <<http://plants.usda.gov>> and Plant Materials Program Web sites <<http://Plant-Materials.nrcs.usda.gov>>.

*The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of*

~~1800/845/4464~~

contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

*To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.*