

# Plant Fact Sheet

## SLENDER HAIRGRASS

## Deschampsia elongata (Hook.) Munro

Plant symbol = DEEL

Contributed by: USDA NRCS Plant Materials Center, Corvallis, Oregon



USDA NRCS Corvallis PMC

#### **Alternate Names**

Another scientific name is Aira elongata.

#### Uses

Slender hairgrass is a moderate to fast growing native bunchgrass useful for erosion control along waterways, ditches, and streambanks, as well as on other seasonally wet or recently disturbed sites. It can also be sown as cover after logging in ravines and other moist forest areas. This species has potential as a nurse crop for slower establishing, longer lived species like California oatgrass (*Danthonia californica*) or other native prairie and

woodland grasses and herbs. Forage value is fair to good prior to maturity. This is a versatile and underutilized native grass. It is also regarded as an attractive ornamental grass because of its fine texture, wispy seed heads, and golden color in summer.

#### Status

Please consult the PLANTS Web site and your State Department of Natural Resources for this plant's current status (e.g. threatened or endangered species, state noxious status, and wetland indicator values).

### **Description and Adaptation**

Slender hairgrass is a moderate sized, fine textured, cool season grass. It is generally a biennial or short-lived perennial under most conditions. Stems (culms) are 40 to 80 (100) cm tall. Leaves are hair-like, less than 1.5 mm wide, and form a basal tuft. The open sheaths are smooth and auricles are absent. The panicle (flowerhead) is 5 to 25 (30) cm long, narrow with slender upward-pointing branches, and a pale green to purple color. This species occurs primarily from Alaska to California and east to Montana and New Mexico.

Key to identification: The slender, upright to arching seed head and fine leaves are indicative of its common name. It should not be confused with tufted hairgrass (Deschampsia caespitosa) which is taller, coarser, and more robust. In addition, the panicles of tufted hairgrass are more open and branching, and the leaves are broader and flatter than slender hairgrass. Annual hairgrass (Deschampsia danthonioides) is smaller and the panicle is also much broader and open compared to slender hairgrass. Because of the fine texture and small florets, it may loosely resemble a bentgrass (Agrostis sp.) without closer inspection.

Relative abundance in the wild: Scattered to relatively common but rarely in large numbers. Slender hairgrass is occasionally found in drainage ways, seep areas, ditches, and damp open woods, as well as adjacent to moist pathways or other disturbed areas.

Adaptation: Slender hairgrass can be found in wet meadows and along the margins of streams, ponds, and lakes. Habitat also includes moist slopes and the understory of open forests from lowland to alpine zones. It is also a component of coastal prairies. As a species of secondary plant succession, slender hairgrass re-establishes after logging in associated forest communities. It is found on fine sandy or

## United States Department of Agriculture-Natural Resources Conservation Service

Plant Materials <a href="http://plant-materials.nrcs.usda.gov/">http://plant-materials.nrcs.usda.gov/</a> Plant Fact Sheet/Guide Coordination Page <a href="http://plant-materials.nrcs.usda.gov/">http://plant-materials.nrcs.usda.gov/</a> intranet/pfs.html> National Plant Data Center <a href="http://ppdc.usda.gov/">http://ppdc.usda.gov/</a>

gravelly soils as well as heavier soil types. This species tolerates full sun to partial shade.

#### **Establishment**

There is no seed dormancy and slender hairgrass establishes readily by sowing in the fall or early spring. Removal of surface hairs (pubescence) on the seed aids in further seed cleaning as well as sowing with a drill. There are approximately 2,300,000 seeds/lb (+/- 20%). Each pound of live seed sown per acre applies about 54 seeds per square foot. Based on seed size and seedling vigor, suggested seeding rates are 2-4 lbs/ac



Line drawing reprinted with permission, University of Washington Press

when sown alone. Higher rates may be required when broadcasting to achieve full stands under more difficult conditions.

### Management

Information on the management of this species is lacking. Given its short-lived nature, grazing must be deferred during flowering and seed set every other year for stand regeneration. Its ability to easily volunteer on moist, disturbed ground and readily produce seed makes it a candidate for moist soil management of wetland habitat for waterfowl, shorebirds, and other wildlife.

## **Environmental Concerns**

Short-lived and may not compete well with weedy herbaceous species. There are no major concerns described for this species.

## Cultivars, Improved, and Selected Materials (and area of origin)

Multiple sources of seed are readily available in California. One or more populations originating from western Oregon are on the market as well. Seed sources from other regions for use in the same area may be lacking.

### Prepared By:

Dale Darris and Robert Tracey, USDA NRCS Plant Materials Center, Corvallis, Oregon.

#### **Species Coordinator:**

Dale Darris, USDA NRCS Plant Materials Center, Corvallis, Oregon.

Edited: 20mar2006 jsp

For more information about this and other plants, please contact your local NRCS field office or Conservation District, and visit the PLANTS Web site<a href="http://plants.usda.gov">http://plants.usda.gov</a> or the Plant Materials Program Web site <a href="http://plant-Materials.nrcs.usda.gov">http://plant-Materials.nrcs.usda.gov</a>

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

Read about <u>Civil Rights at the Natural Resources Convervation</u> Service.