



## EASTERN HEMLOCK

*Tsuga canadensis* (L.) Carr.

plant symbol = TSCA

Contributed By: USDA, NRCS, National Plant Data Center & the Biota of North America Program



© R.A. Seelig  
Botany Dept., NMNH, Smithsonian Institution  
@ PLANTS

### Alternate common names

Canada hemlock, hemlock spruce

### Uses

Eastern hemlock was early valued for its bark, an important source of tannin for the leather industry. Trees were felled and stripped of their bark, which was then milled for tannin extraction. To simplify stripping the bark and turning the logs, trees were often felled into lakes. Many of these logs were much later extracted from northern lakes and milled.

The wood has been used for light framing, roofing, boxes and crates, and pulping, but it tends to be brittle and eastern hemlock is not presently important as a timber tree. Commercial stands have been greatly reduced by prior harvesting and lack of restocking.

American Indians used the cambium as the base for breads and soups or mixed it with dried fruit and animal fat for pemmican. Natives and white settlers also made tea from hemlock leaves, which have a high vitamin C content.

Eastern hemlock can be used as a specimen, screen, or group planting, and it can be sheared over time into a formal evergreen hedge, which is densely leafy all the way to the ground (due to its full shade

tolerance), although as a hedge it must be repeatedly pruned to keep it in size. It has a naturally open growth habit – if bought with a dense canopy effect, it may have been repeatedly sheared at the nursery (or Christmas tree farm).

Numerous cultivars of eastern hemlock have been developed, including compact shrubs, dwarfs, form mutants (weeping, fastigiate, prostrate, etc.), color mutants (variegated), and graceful trees. The wild type apparently also is common in cultivation. Eastern hemlock stands are considered important as shelter and cover for white-tailed deer and other wildlife species, such as turkey, ruffed grouse, and others.

### Status

Please consult the PLANTS Web site and your State Department of Natural Resources for this plant's current status, such as, state noxious status and wetland indicator values.

### Description

*General:* Pine family (Pinaceae). Native trees to 30 meters tall, with a broadly conic crown, the branches often drooping at the ends and “feathery;” twigs yellow-brown, densely pubescent. Bark brownish, scaly and fissured. Needles evergreen, flat, mostly appearing 2-ranked, (5-)15-20(-25) mm long, narrowed to a petiole-like base and set on peg-like projections, the lower surface waxy, with 2 broad, conspicuous stomatal bands (like pale lines), the upper surface shiny-green to yellow-green, the margins minutely toothed, especially toward apex. Seed cones ovoid, 1.5-2.5 cm long, borne near the branch tips, hanging. The common name pertains to its distribution in eastern North America.

*Variation within the species:* various studies have recorded physiological and morphological variation within the species, but no major discontinuities have been found to suggest that named entities should be recognized.

The geographic distribution of eastern hemlock completely overlaps that of Carolina hemlock (*Tsuga caroliniana*), which differs in its high elevation habitats, leaves mostly spreading all directions from the twigs, and more elongate seed cones.

*Distribution:* Eastern hemlock is a species of the northeastern and Appalachian regions of North America: from New Brunswick and Nova Scotia to southern Quebec and Ontario, south to northern

Georgia and Alabama, west in the lake states to Minnesota with outliers in southern Michigan, western Ohio, and southern Indiana. Many disjunct populations, probably glacial relicts, occur east of the Appalachians in the middle Atlantic states. For current distribution, please consult the Plant Profile page for this species on the PLANTS Web site.

### **Adaptation**

Eastern hemlock grows from about sea level to 1500 meters. In the more southern parts of its range, eastern hemlock occurs only where there is drainage of cool, moist air -- on moist rocky ridges, valleys and ravines, hillsides, and lakeshores. In the northern hardwood forest, it is found on a greater variety of sites, including low rolling hills and glacial ridges. Eastern hemlock most commonly grows in mixed stands with species such as white pine, northern red oak, sugar maple, American beech, yellow birch, and white ash.

### **Establishment**

Eastern hemlocks begin to produce cones at about age 15 in vigorous trees or much later in suppressed trees. There is a high frequency of cone crops and individual trees have a long duration of cone production (excellent cone production has been reported for trees more than 450 years old), but the viability of seed usually is low. Seeds are particularly easily damaged by drying. The best conditions for germination and seedling establishment are under a 70-80% crown cover on an exposed, partially decomposed layer. Otherwise, regeneration is restricted to rotten logs, stumps, and mounds that normally have warmer surfaces and better moisture retention than the forest floor.

Eastern hemlock is the most shade tolerant of all tree species and individuals may remain in the understory in natural stands for 25-400 years. After a plant reaches 1-2 meters in height, the root system has reached a depth not radically affected by surface drying, and release from overstory competition may greatly increase annual growth. In general, eastern hemlock is a slow-growing tree that may take 250 to 300 years to reach maturity and may live for 900 years or more.

### **Management**

Eastern hemlock generally does not tolerate nutrient-poor soils, wet soils or poorly drained sites, prolonged drought, prolonged heat, sun scorch, windy and exposed sites, aerial pollution, or winter salt spray. Drought is probably the most serious damaging agent to the species, especially during the seedling stage. Damping-off fungi and root rots also

are seriously damaging to young plants. Containerized plants are best for transplanting -- move into sites that are cool, well-drained, and wind-protected, in partial sun to partial shade. Good drainage is essential for transplant success -- the porosity of the soil should be improved with peat moss or sand, with the root ball elevated about 2" to 4" above the surrounding soil grade.

A shallow root system makes trees highly susceptible to wind-throw when exposed through timber cutting or planted in open sites. Plants should be staked for the first two or three years following transplant, to prevent wind-throw. Saplings and small trees are highly susceptible to damage from fire because of the thin bark, and root injury often occurs from high intensity fires because of heavy litter concentration.

The most severe insect pest is the Asian hemlock woolly adelgid, a phloem-feeding insect that causes branch dieback and tree decline. Trees typically die after several years of adelgid infestation.

### **Cultivars, Improved and Selected Materials (and area of origin)**

Please check the Vendor Database, expected to be on-line through the PLANTS Web site in 2001 by clicking on Plant Materials. These plant materials are readily available from commercial sources.

### **References**

Godman, R.M. & K. Lancaster 1990. *Tsuga canadensis*. Pp. 604-612, IN: R.M. Burns and B.H. Honkala. *Silvics of North America. Volume 1. Conifers*. USDA Forest Service Agric. Handbook 654, Washington, D.C.  
<[http://willow.ncfes.umn.edu/silvics\\_manual/Table\\_of\\_contents.htm](http://willow.ncfes.umn.edu/silvics_manual/Table_of_contents.htm)>

Taylor, R.J. 1993. *Tsuga*. Pp. 362-365, IN: *Flora of North America, north of Mexico*. Vol. 2, *Pteridophytes and Gymnosperms*. Oxford Univ. Press, New York. <[http://hua.huh.harvard.edu/cgi-bin/Flora/flora.pl?FLORA\\_ID=12395](http://hua.huh.harvard.edu/cgi-bin/Flora/flora.pl?FLORA_ID=12395)>

USDA, NRCS 2000. *The PLANTS database*. 21SEP2000. National Plant Data Center, Baton Rouge, Louisiana. <<http://plants.usda.gov>>

### **Prepared By**

*Guy Nesom*  
BONAP, North Carolina Botanical Garden,  
University of North Carolina, Chapel Hill, North Carolina

**Species Coordinator**

*Lincoln Moore*

USDA, NRCS, National Plant Data Center, Baton Rouge, Louisiana

Edited 17jan01 jsp

For more information about this and other plants, please contact your local NRCS field office or Conservation District, and visit the PLANTS <<http://plants.usda.gov>> and Plant Materials Program Web sites <<http://Plant-Materials.nrcs.usda.gov>>.

*The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).*

*To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.*