

GREEN SILKYSCALE

Anthraenantia villosa (Michx.)

Beauv.

Plant Symbol = ANVI4

Contributed By: USDA NRCS National Plant Data Center



From Hitchcock, A.S. (1950)
@ plants.usda.gov

Alternate Name

green silkyscale

Uses

Green silkyscale is seldom abundant enough to be a key management species but it adds variety to livestock diet. Cattle graze it readily during summer, fall, and early winter. It is an indicator of good range condition. Adapts to well-drained upland soils.

Status

Please consult the PLANTS Web site and your State Department of Natural Resources for this plant's

current status, such as, state noxious status and wetland indicator values.

Description

Green silkyscale is a native, warm-season, rhizomatous perennial grass. The height is between 2 and 4 feet. The leaf blade is 4 to 12 inches long; light green; a distinct fringe of silky hair along edge; and twists and curls particularly when young. The leaf sheath is chiefly basal, rounded, and with an obscure collar. The ligule is ridge-like membrane with a dense ring of short hair. The stem is round and slender. The seedhead is a slender panicle, pale green with oval spikelets covered with short, stiff hairs.

Distribution: For current distribution, please consult the Plant Profile page for this species on the PLANTS Web site.

Management

This grass decreases when closely grazed throughout its growing season. To maintain this species in the plant community, defer grazing 70 to 90 days every 3 to 4 years during the growing season. Foliage can be burned if done during the dormant season and no more than every 2 to 3 years. Grazing should be deferred 90 days during the spring growing season following a burn.

Establishment

Green silkyscale growth starts slowly in spring and becomes more prominent during summer. It forms loose colonies, forms seedheads in the fall, and becomes dormant in the fall and winter. The seedstalks break off and disappear in the winter. It grows throughout the South from the 30-inch rainfall belt eastward and is adapted to well-drained upland soils.

Cultivars, Improved and Selected Materials (and area of origin)

Please contact your local NRCS Field Office.

Reference

Leithead, H.L., L.L. Yarlett, & T.N. Shiflett. 1976. 100 native forage grasses in 11 southern states. USDA SCS *Agriculture Handbook No. 389*, Washington, DC.

Prepared By & Species Coordinator:

Percy Magee

USDA NRCS National Plant Data Center, Baton
Rouge, Louisiana

Edited: 13may02 ahv; jul03 ahv; 20sep05 jsp; 070112 jsp

For more information about this and other plants, please contact your local NRCS field office or Conservation District, and visit the PLANTS Web site <<http://plants.usda.gov>> or the Plant Materials Program Web site <<http://Plant-Materials.nrcs.usda.gov>>

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's [TARGET Center](#) at 202-720-2600 (voice and TDD).

To file a complaint of discrimination write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

Read about [Civil Rights at the Natural Resources Conservation Service](#).