

## EVERGREEN HUCKLEBERRY

*Vaccinium ovatum* Pursh  
Plant Symbol = VAOV2

Contributed by: USDA NRCS National Plant Data Center & Oregon Plant Materials Center



Alfred Brousseau  
© Brother Eric Vogel, St. Mary's College  
@ CalPhotos

### Alternate Names

California huckleberry, shot huckleberry, huckleberry, winter huckleberry, evergreen huckleberry

### Uses

*Ethnobotanic:* Tribes in British Columbia and western Washington use the berries of evergreen huckleberry. These tribes include the Sechelt, Comox, Straits, Halkomelem, Lower Nlaka'pamux Salish, Nuu-chah-nulth (Vancouver Island's West Coast), and the Quinault of Washington. Evergreen huckleberries were well liked and people often traveled great distances to obtain them. The berries ripen late in the year, around October or November. They are the last fruits to be gathered in the season round and are said to be even tastier after freezing. The berries are eaten fresh, usually with oil. The berries are also sun or smoke dried, partly mashed, pressed into cake form, and wrapped in leaves or bark. Today they are made into jam or used in cooking.

The leaves and berries are high in vitamin C. The leaves and finely chopped stems contain quinic acid, a former therapeutic for gout said to inhibit uric acid formation but never widely used because of mixed clinical results. The leaves have been widely used to lower or modify blood sugar levels. Many herbalists maintain that huckleberry leaf tea may be useful in

stabilizing blood sugar levels in cases of diabetes, and medical research has shown that consumption of the leaf extract decreases blood sugar levels shortly after administration. Taken on regular basis, huckleberry tea will gradually help alleviate both glycosuria and hyperglycemia and appears to have a beginning, but useful effect as an adjunct treatment to diabetes mellitus. The leaves are believed also to stimulate appetite, and have astringent and antiseptic qualities that are useful in urinary disorders.

*Horticulture:* Evergreen huckleberry is an excellent horticultural choice due to its beautiful, glossy, evergreen foliage and tolerance of a wide range of light levels. The foliage is often used in flower arrangements.

*Wildlife & Livestock:* The foliage of evergreen huckleberry is browsed by elk and deer. Flowers attract butterflies. For several species of grouse, huckleberries are among the most important summer and early fall foods. Berries are eaten by chipmunks, black bear, mice, scarlet tanagers, bluebirds, thrushes, and other songbirds. Deer and rabbit browse freely on the plants. Because of their food value to wildlife and their dense shrubby growth, evergreen huckleberry is worthy of inclusion in hedgerows.

In some localities goats and deer crop evergreen huckleberry rather closely, utilizing 30 to 40 % of the leafage and current twigs. Sheep crop it somewhat less closely but it enters into their diet to a considerable extent in late summer and autumn. The browse rating is fair to poor for sheep, goats, and deer; poor to useless to cattle; and useless for horses.

### Status

Please consult the PLANTS Web site and your State Department of Natural Resources for this plant's current status, such as, state noxious status and wetland indicator values.

### Description

*General:* Heath Family (Ericaceae). This erect, evergreen shrub is stout, from 0.5-3 m tall. The glossy green leaf blades are 2-5 cm, ovate, leathery, serrate, with glandular hairs on the lower surface. The umbel-like inflorescence emerges from the leaf axils. Urn-shaped flowers are bright pink. The berries are 6-9 mm, purplish-black. Evergreen huckleberry does not generally root easily.

## Distribution

For current distribution, please consult the Plant Profile page for this species on the PLANTS Web site. Evergreen huckleberry grows from the west side of the Cascades in Washington to the coast of British Columbia, to the redwood area of California. It is sporadic south to Santa Barbara, California and in the coast ranges to the central Sierra Nevada Mountains.

## Establishment

**Adaptation:** *Vaccinium ovatum* grows in edges and clearings of coniferous woods, at elevations from 3-800 m. Evergreen huckleberry can also be found near beaches in the salt spray zone. This huckleberry grows in moist to slightly dry soils. It will grow in full sun to full shade, although the plants prefer some shade.



Jeanne Russell Janish  
Used with permission of the publishers  
Stanford University

**Propagation:** Evergreen huckleberry can be difficult to propagate or transplant, but it is available in some nurseries. It can be grown from cuttings, from seed, or by layering. Huckleberry cuttings should be taken while the plant is dormant, from November to April. Their rooting success is fairly sporadic.

Evergreen huckleberry requires excessive drainage and acidic soils to become established. It does best in full or partial shade; it may tolerate morning and winter sun.

**Live Plant Collections:** Evergreen huckleberry is propagated by cuttings from fully matured shoots taken in fall and winter, when the plant is dormant. Cuttings made from the previous year's growth taken the third week in April rooted 100% (Vancouver, B.C.). Application of 0.3 to 0.4% IBA talc to the freshly cut stem surface and basal heat (21°C; 70° F) to potted plants will enhance rooting.

Young plants can be salvaged, but they should be transplanted when they are less than one foot tall. Frequently, these small plants will turn out to be new shoots of a mature plant reviving from deer browsing or logging, and will die from lack of roots.

**Seed Collections:** Berries should be collected when they are ripe (from August to September or later). The blue-black fruit is easily collected by hand picking or by beating the bush over a large bucket. Following collection, chill the fruit at 10°C for several days. Clean seeds by macerating and floating off the pulp and unsound seed. Clean seeds carefully; they are minuscule, so you may want to use pantyhose or cheesecloth to strain the seed from the pulp.

Seeds dried at 15-21°C for two days can be stored in a refrigerator for up to 12 years. Fresh seeds not planted in the fall may germinate better if cold stratified for 1-3 months. Stored seeds germinates well when exposed to alternating temperature and light regimes of 28°C light for 14 hours a day and 13°C dark for 10 hours.

Fresh or stored and cold-stratified seeds can be sown directly into flats or small pots (a salt shaker can be used for sowing). Plant in a mixture of sand and peat moss. Seedlings will begin to emerge in a month and will continue to emerge for a long period thereafter. Transplant seedlings into larger pots 6 to 7 weeks after emergence. Plant outside after the first growing season. Seedlings are slow growing, and it may take 2-3 years for a nursery-sized plant to develop.

## Management

This plant grows very rapidly in moist, shady conditions. If summer drought occurs, the plants should be watered so roots are kept fairly moist.

**Traditional Resource Management:** This includes the following: 1) occasional burning to stimulate new growth; 2) pruning the branches after picking the berries to stimulate new growth and fruit production the next growing season; and 3) ownership of red huckleberry shrubs provides the basis for careful tending and sustainable yield of valued resources.

## Cultivars, Improved and Selected Materials (and area of origin)

This species is readily available from native plant nurseries within its range. Contact your local Natural Resources Conservation Service (formerly Soil Conservation Service) office for more information. Look in the phone book under "United States Government." The Natural Resources Conservation

Service will be listed under the subheading "Department of Agriculture."

## References

Abrams, L. & R.S. Ferris 1960. *Illustrated flora of the Pacific states*. 4 Vols. Stanford University Press, Palo Alto, California.

Barton, B.J. 1997. *Gardening by mail. A source book*. A Mariner Book. Houghton Mifflin Company, Boston and New York.

Cooke, S.S. 1997. *A field guide to the common wetland plants of western Washington and northwestern Oregon*. Seattle Audubon Society and Washington Native Plant Society. 414 pp.

Crossley, J.A. 1974. *Vaccinium L. blueberry*. pp. 840-843 IN: Shopmeyer, C.S. (tech. coord.) 1974. *Seeds of the Woody Plants in the United States*. Agric. Handbook 450. Washington, D.C. USDA, Forest Service. 883 pp.

Gunther, E. 1973. *Ethnobotany of western Washington*. University of Washington Publications in Anthropology, 10(1). University of Washington Press, Seattle, Washington.

Harrington, H.D. 1972. *Western edible wild plants*. The University of New Mexico Press. 156 pp.

Hartmann, H.T., D.E. Kester, & F.T. Davies, Jr. 1990. *Plant propagation principles and practices*. Prentice Hall, Englewood Cliffs, New Jersey.

Hitchcock, C.L. & A. Cronquist 1973. *Pacific Northwest flora*. University of Washington Press, Seattle and London. 730 pp.

Isaacson, R.T. 1993. *Anderson horticultural library's source list of plants and seeds*. Anderson Horticultural Library. University of Minnesota Libraries. Minnesota Landscape Arboretum. 261 pp.

Kunlein, H.V. & N.J. Turner 1991. *Traditional plant foods of Canadian indigenous peoples. Nutrition, botany, and use*. Food and Nutrition in History and Anthropology Volume 8. Gordon and Breach Science Publishers. 632 pp.

Leigh, M. (August) 1997. *Grow your own native landscape: A guide to identifying, propagating, and landscaping with western Washington native plants*. Environmental Protection Agency, The Washington State Department of Ecology, and Washington State University Cooperative Extension.

Martin, A.C., H.S. Zim, & A.L. Nelson 1951. *American wildlife and plants: A guide to wildlife food habits*. Dover Publications, Inc., New York, New York. 500 pp.

Moore, M. 1979. *Medicinal plants of the mountain west*. Museum of New Mexico Press. 200 pp.

Rose, R., C.E.C. Chachulski, & D. Haase 1998. *Propagation of Pacific Northwest native plants*. Oregon State University Press, Corvallis, Oregon.

Sampson, A.W. & B.S. Jespersen 1981. *California range brushlands and browse plants*. Division of Agricultural Sciences, University of California, Berkeley, California. 162 pp.

Schlick, M.D. 1994. *Columbia River basketry. Gift of the ancestors, gift of the Earth*. University of Washington Press, Seattle and London. 232 pp.

Schopmeyer, C.S. (Tech. Coord.) 1974. *Seeds of woody plants in the United States*. Agriculture Handbook No. 450. USDA, Forest Service, Washington, D.C.

Stevens, M. & R. Vanbianchi 1993. *Restoring wetlands in Washington. A guidebook for wetland restoration, planning and implementation*. Washington State Department of Ecology. Publication #93-17.

Turner, N.J., L.C. Thompson, M.T. Thompson & A.Z. York 1990. *Thompson ethnobotany: Knowledge and usage of plants by the Thompson Indians of British Columbia*. Royal British Columbia Museum Memoirs No. 3, Victoria, British Columbia, Canada

Turner, N.J., J. Thomas, B.F. Carlson & R.T. Ogilvie 1983. *Ethnobotany of the Nitinaht Indians of Vancouver Island*. British Columbia Provincial Museum, Occasional Paper No. 24. 165 pp.

Turner, N.J. & B.S. Efrat 1982. *Ethnobotany of the Hesquiat Indians of Vancouver Island*. B.C. Provincial Museum Cultural Recovery Paper No. 2. 99 pp.

Turner, N.J. 1978. *Food plant of British Columbia Indians. Part II. Interior peoples*. British Columbia Provincial Museum Handbook No. 36, Victoria, British Columbia, Canada.

Turner, N.J. 1975. *Food plants of British Columbia Indians. Part I. Coastal Peoples*. British Columbia

Provincial Museum Handbook No. 34, Victoria,  
British Columbia, Canada.

Vanbianchi, R., M. Stevens, T. Sullivan & S.  
Hashisaki 1994. *A citizen's guide to wetland  
restoration*. U.S. Environmental Protection Agency,  
Region 10. 71 pp.

**Prepared By**

*Michelle Stevens*

Formerly USDA, NRCS, National Plant Data Center

*Dale C. Darris*

USDA, NRCS, Plant Materials Center, Corvallis,  
Oregon

**Species Coordinator**

*M. Kat Anderson*

USDA, NRCS, National Plant Data Center  
c/o Plant Sciences Department, University of  
California, Davis, California

Edited: 17jan01 jsp; 060818 jsp

For more information about this and other plants, please contact  
your local NRCS field office or Conservation District, and visit the  
PLANTS Web site <<http://plants.usda.gov>> or the Plant Materials  
Program Web site <<http://Plant-Materials.nrcs.usda.gov>>

*The U.S. Department of Agriculture (USDA) prohibits  
discrimination in all its programs and activities on the basis of  
race, color, national origin, sex, religion, age, disability, political  
beliefs, sexual orientation, and marital or family status. (Not all  
prohibited bases apply to all programs.) Persons with disabilities  
who require alternative means for communication of program  
information (Braille, large print, audiotape, etc.) should contact  
USDA's [TARGET Center](#) at 202-720-2600 (voice and TDD).*

*To file a complaint of discrimination write USDA, Director, Office  
of Civil Rights, Room 326-W, Whitten Building, 14th and  
Independence Avenue, SW, Washington, DC 20250-9410 or call  
202-720-5964 (voice or TDD). USDA is an equal opportunity  
provider and employer.*

*Read about [Civil Rights at the Natural Resources Conservation  
Service](#).*